

# COMMUNICATIONS

DE LA FACULTÉ DES SCIENCES  
DE L'UNIVERSITÉ D'ANKARA

Série A<sub>1</sub>: Mathématiques

---

TOME 29

ANNÉE 1980

---

**The Space of Sequences of Which A-Transforms Are In  $l_p$**

by

**Öner ÇAKAR**

2

Faculté des Sciences de l'Université d'Ankara  
Ankara, Turquie

# Communications de la Faculté des Sciences de l'Université d'Ankara

Comité de Rédaction de la Série A<sub>1</sub>

B. Yurtsever H. Hacısalihođlu M. Oruç

Secrétaire de publication.

A. Yalçın

La Revue "Communications de la Faculté des Sciences de l'Université d'Ankara" est un organe de publication englobant toutes les disciplines scientifiques représentées à la Faculté.

La Revue, jusqu'à 1975 à l'exception des tomes I, II, III, était composée de trois séries:

Série A: Mathématiques, Physique et Astronomie.

Série B: Chimie.

Série C: Sciences naturelles.

A partir de 1975 la Revue comprend sept séries:

Série A<sub>1</sub>: Mathématiques

Série A<sub>2</sub>: Physique

Série A<sub>3</sub>: Astronomie

Série B : Chimie

Série C<sub>1</sub>: Géologie

Série C<sub>2</sub>: Botanique

Série C<sub>3</sub>: Zoologie

En principe, la Revue est réservée aux mémoires originaux des membres de la Faculté. Elle accepte cependant, dans la mesure de la place disponible, les communications des auteurs étrangers. Les langues allemande, anglaise et française sont admises indifféremment. Les articles devront être accompagnés d'un bref sommaire en langue turque.

Adres: Fen Fakültesi Tebliğler Dergisi Fen Fakültesi, Ankara, Turquie.

# The Space of Sequences of Which A-Transforms Are In $l_p$

Öner ÇAKAR

(University of Ankara, Faculty of Science Dept. of Mathematics)

(Received on 21 December, 1979 and accepted on 3 March, 1980)

## SUMMARY

In this paper, we have defined and investigated two new sequence spaces  $l_p(A)$  and  $l_\infty(A)$  which contain the sequence spaces  $X_{a(p)}$  and  $X_{a(\infty)}$ , given by Wang, [5], as special cases, respectively. Some inclusion theorems between the related sequence spaces have also been proved.

## 1. Introduction.

In [5], Wang has defined Nörlund sequence spaces  $X_{a(p)}$  and  $X_{a(\infty)}$  as

$$X_{a(p)} = \left\{ x = (x_k) : \left( \sum_{n=0}^{\infty} \left| \frac{1}{A_n} \sum_{k=0}^n a_{n-k} x_k \right|^p \right)^{1/p} < \infty, 1 \leq p < \infty \right\}$$

and

$$X_{a(\infty)} = \left\{ x = (x_k) : \sup_n \left| \frac{1}{A_n} \sum_{k=0}^n a_{n-k} x_k \right| < \infty, n \geq 0 \right\}$$

respectively, where  $a = (a_n)$  is a sequence of positive real numbers

and  $A_n = \sum_{k=0}^n a_k$ . In other words, the spaces  $X_{a(p)}$  and  $X_{a(\infty)}$

consist of the sequences of which Nörlund transforms are in  $l_p$  and in  $l_\infty$ , respectively, where

$$l_p = \{ x = (x_k) : \sum_{k=1}^{\infty} |x_k|^p < \infty, 1 \leq p < \infty \}$$

and

$$l_\infty = \{ x = (x_k) : \sup_k |x_k| < \infty \}.$$

It is known that the sequence space

$$X_p = \{ x = (x_k) : (\sum_{n=1}^{\infty} | \frac{1}{n} \sum_{k=1}^n x_k |^p)^{1/p} < \infty, 1 \leq p < \infty \}$$

which has been introduced by Ng, [2], is a special case of  $X_{a(p)}$ , corresponding to  $a_n = 1$  for every  $n$ , [5]. Obviously, the space  $X_p$  consists of the sequences of which (C,1)-transforms are in  $l_p$ .

We note that if  $A = (a_{nk})$  is an infinite matrix of numbers  $a_{nk}$  ( $n, k = 1, 2, \dots$ ) then the sequence  $(A_n(x))$  given by the matrix multiplication

$$A_n(x) = \sum_{k=1}^{\infty} a_{nk} x_k$$

(assuming that  $A_n(x)$  exists for every  $n = 1, 2, \dots$ ) is called the  $A$ -transform of the sequence  $x = (x_k)$ .

In this article, we will first define two new sequence spaces  $l_p(A)$  and  $l_{\infty}(A)$  which consist of the sequences of which any  $A$ -transform is in  $l_p$  and in  $l_{\infty}$ , respectively. And then, we will investigate some properties of these spaces.

Throughout the paper, the triangular inequality, Minkowski's inequality, Hölder's inequality and the following inequality ([1], p.4)

$$(1) \quad \left( \sum_{k=1}^n |a_k|^s \right)^{1/s} \leq \left( \sum_{k=1}^n |a_k|^r \right)^{1/r}, \quad (0 < r < s)$$

will be used frequently.

## 2. Definitions.

Let  $x = (x_k)$  be a sequence of real numbers and  $A = (a_{nk})$  be an infinite matrix of positive real numbers. Now, let us define

$$(2) \quad l_p(A) = \{ x = (x_k) : \sum_{n=1}^{\infty} | \sum_{k=1}^{\infty} a_{nk} x_k |^p < \infty, 1 \leq p < \infty \}$$

and

$$(3) \quad l_{\infty}(A) = \{ x = (x_k) : \sup_n | \sum_{k=1}^{\infty} a_{nk} x_k | < \infty, n \geq 1 \}.$$

It can easily be seen that when we take the identity matrix, (C,1) matrix and  $N_a$  matrix for  $A = (a_{nk})$  in  $l_p(A)$ , we get the sequence spaces  $l_p$ ,  $X_p$  and  $X_{a(p)}$  as the special cases, respectively.

After some routine calculations it becomes clear that the spaces  $l_p(A)$  and  $l_\infty(A)$  are Banach spaces with the norms

$$\|x\|_p = \left( \sum_{n=1}^{\infty} \left| \sum_{k=1}^{\infty} a_{nk} x_k \right|^p \right)^{1/p}, \quad (1 \leq p < \infty)$$

and

$$\|x\|_\infty = \sup_n \left| \sum_{k=1}^{\infty} a_{nk} x_k \right|, \quad (n \geq 1)$$

respectively.

### 3. Inclusion Theorems

In this paragraph, we are going to give some inclusion theorems between the related sequence spaces.

**Theorem 1.** If  $1 \leq r < s$  then  $l_r(A) \subseteq l_s(A)$ .

**Proof.** The proof of this theorem is an immediate consequence of the inequality (1).

**Theorem 2.** Let  $A = (a_{nk})$  be a triangular matrix of positive numbers and  $x = (x_k)$  be a sequence of real numbers. If

$$(4) \quad a_{nn} = O(n^{-1})$$

and  $(a_{nk})_{1 \leq k \leq n}$  forms a non-decreasing sequence for each  $n$  then

$$ces_p \subseteq l_p(A)$$

where  $ces_p$  is the sequence space given by

$$ces_p = \left\{ x = (x_k) : \left( \sum_{n=1}^{\infty} \left( \frac{1}{n} \sum_{k=1}^n |x_k| \right)^p \right)^{1/p} < \infty, 1 < p < \infty \right\}$$

([4]).

**Proof.** Let  $x = (x_k) \in ces_p$ . Using the triangular inequality and condition (4), we write

$$\left| \sum_{k=1}^n a_{nk} x_k \right|^p \leq \left( \sum_{k=1}^n a_{nk} |x_k| \right)^p$$

$$\begin{aligned} &\leq (a_{nn} \sum_{k=1}^n |x_k|)^p \\ &= O(1) \left( \frac{1}{n} \sum_{k=1}^n |x_k| \right)^p \end{aligned}$$

Then, taking sum over  $n$  from 1 to  $\infty$ , we get

$$\sum_{n=1}^{\infty} \left| \sum_{k=1}^n a_{nk} x_k \right|^p \leq O(1) \sum_{n=1}^{\infty} \left( \frac{1}{n} \sum_{k=1}^n |x_k| \right)^p < \infty$$

which completes the proof.

**Theorem 3.** Let  $A = (a_{nk})$  and  $x = (x_k)$  be as in Theorem 2. If

$$(5) \quad a_{nn} = O(n^{-1})$$

and  $(a_{nk})_{1 \leq k \leq n}$  forms a non-increasing sequence for each  $n$ , then

$$\text{ces}_p \subseteq I_p(A).$$

**Proof.** Let  $x = (x_k) \in \text{ces}_p$ . Then using the triangular inequality and condition (5), we write

$$\begin{aligned} \left| \sum_{k=1}^n a_{nk} x_k \right|^p &\leq \left( \sum_{k=1}^n a_{nk} |x_k| \right)^p \\ &\leq (a_{n1} \sum_{k=1}^n |x_k|)^p \\ &= O(1) \left( \frac{1}{n} \sum_{k=1}^n |x_k| \right)^p \end{aligned}$$

and so we get

$$\sum_{n=1}^{\infty} \left| \sum_{k=1}^n a_{nk} x_k \right|^p \leq O(1) \sum_{n=1}^{\infty} \left( \frac{1}{n} \sum_{k=1}^n |x_k| \right)^p < \infty$$

which proves the theorem.

**Theorem 4.** Let  $A = (a_{nk})$  and  $B = (b_{nk})$  be any two normal infinite matrices of positive real numbers, ([3], pp. 14–15). And let  $C = (c_{nk})$  be a matrix defined by  $C = BA^{-1}$  satisfying the condition

$$(6) \quad \sum_{k=1}^n |c_{nk}| = O(n^{-1})$$

where  $A^{-1}$  is the inverse matrix of  $A$ , then

$$l_p(A) \subseteq l_p(B), \quad p > 1.$$

**Proof.** Let  $x = (x_k) \in l_p(A)$ . Since

$$b_{nk} = \sum_{i=1}^n c_{ni} a_{ik}$$

by the definition of the matrix  $C$ , we write

$$\begin{aligned} \sum_{k=1}^n b_{nk} x_k &= \sum_{k=1}^n \left( \sum_{i=1}^n c_{ni} a_{ik} \right) x_k \\ &= \sum_{k=1}^n c_{nk} \sum_{i=1}^k a_{ki} x_i \end{aligned}$$

Put

$$M = \sum_{k=1}^{\infty} \left| \sum_{i=1}^k a_{ki} x_i \right|^p.$$

Then using Hölder's inequality and condition (6), we have

$$\begin{aligned} \left| \sum_{k=1}^n b_{nk} x_k \right| &= \left| \sum_{k=1}^n c_{nk} \sum_{i=1}^k a_{ki} x_i \right| \\ &\leq \left( \sum_{k=1}^n |c_{nk}|^q \right)^{1/q} \left( \sum_{k=1}^n \sum_{i=1}^k |a_{ki} x_i|^p \right)^{1/p} \\ &\leq M^{1/p} \sum_{k=1}^n |c_{nk}|, \end{aligned}$$

where  $p^{-1} + q^{-1} = 1$ . Therefore, we finally get

$$\sum_{n=1}^{\infty} \left| \sum_{k=1}^n b_{nk} x_k \right|^p \leq O(1) M \sum_{n=1}^{\infty} \frac{1}{n^p} < \infty$$

which completes the proof.

#### ÖZET

Bu çalışmada, Wang tarafından [5] de verilen  $X_{a(p)}$  ve  $X_{a(\infty)}$  dizi uzaylarını özel hal olarak içeren  $l_p(A)$  ve  $l_{\infty}(A)$  dizi uzayları tanımlanarak, bu uzayların temel özellikleri incelenmiş ve ilgili dizi uzayları arasındaki bazı içermeye bağıntıları ispatlanmıştır.

## REFERENCES

- [1] G.H.HARDY, J.LITTLEWOOD and G.POLYA, *Inequalities*, Cambridge University Press, 1967.
- [2] NG PENG-NUNG, On modular sequence space of a non-absolute type, *Nanta Math.*, 2 (1978) 84-93.
- [3] A.PEYERIMHOFF, *Lectures on Summability*, Lecture Notes in Mathematics, Springer-Verlag, Vol. 107,1969.
- [4] JAU-SHYONG SHIUE, On the Cesàro sequence spaces, *Tamkang Journal of Mathematics*, 1 (1970) 19-25.
- [5] CHUG-SHIN WANG, On Nörlund sequence spaces, *Tamkang Journal of Mathematics*, 9 (1978) 269-274.



**Prix de l'abonnement annuel**

**Turquie: 15 TL; Etranger: 30 TL.**

**Prix de ce num ero: 5 TL (pour la vente en Turquie).**

**Pri ere de s'adresser pour l'abonnement  : Fen Fak ltesi  
Dekanlıđı Ankara, Turquie.**