

CURVATURE OF SPHERES IN MINKOWSKI (n+1)-SPACE

E.M. ELKHOLY

Mathematics Department, Tanta University - Egypt.

(Received: 4 February 1987. Accepted: 20 May 1987)

ABSTRACT

In this paper we show that the hypersurfaces $S(\alpha)$ in Minkowski $(n+1)$ -space M^{n+1} given by:

$$x_1^2 + \dots + x_n^2 + \alpha = x_{n+1}^2, \quad \alpha \neq 0,$$

carry induced Riemannian metrics which is positive definite in case $\alpha > 0$ and negative definite in case $\alpha < 0$. Also the curvature of $S(\alpha)$ is constant and equal to $-n(n-1)/\alpha$ for $\alpha > 0$ and $n(n-1)/\alpha$ for $\alpha < 0$.

THE MAIN RESULT

The "positive spheres" $S(\alpha)$ in Minkowski $(n+1)$ -space M^{n+1} is given by:

$$\{(x_1, x_2, \dots, x_{n+1}) \in \mathbb{R}^{n+1} : x_1^2 + \dots + x_n^2 + \alpha = x_{n+1}^2, \alpha \neq 0\}$$

This is a hyperboloid of revolution of two sheets diffeomorphic to $\mathbb{R}^n \times S^0$. Thus $S(\alpha)$ is a smooth hypersurface in \mathbb{R}^{n+1} . The Lorentz structure on \mathbb{R}^{n+1} induces a symmetric bilinear form on the tangent space $T_x S$ to S at each $x \in S$. This form h can be calculated as follows:

Without loss of generality we can restrict attention to the positive spheres S_+ where $x_{n+1} > 0$, and $\alpha > 0$. There is a coordinate system on S_+ given by projection to the first n coordinates in \mathbb{R}^{n+1} , and

$$ds^2 = dx_1^2 + dx_2^2 + \dots + dx_n^2 - dx_{n+1}^2, \text{ but}$$

$$x_{n+1}^2 = x_1^2 + x_2^2 + \dots + x_n^2 + \alpha, \text{ and hence}$$

$$dx_{n+1}^2 = (x_1 dx_1 + \dots + x_n dx_n)^2 / \Delta,$$

where

$$\Delta = x_1^2 + \dots + x_n^2 + \alpha.$$

Now ds^2 takes the form

$$\begin{aligned} ds^2 &= \sum_{k=1}^n dx_k^2 - \frac{1}{\Delta} \sum_{i,j=1}^n x_i x_j dx_i dx_j \\ &= \sum_{i,j=1}^n h_{ij} dx_j, \end{aligned}$$

thus the matrix H_x representing the form h_x for any $x \in S$ is given by

$$H_x = \begin{bmatrix} 1 - \frac{x_1^2}{\Delta} & -\frac{x_1 x_2}{\Delta} & \dots & -\frac{x_1 x_n}{\Delta} \\ -\frac{x_1 x_2}{\Delta} & 1 - \frac{x_2^2}{\Delta} & \dots & -\frac{x_2 x_n}{\Delta} \\ \dots & \dots & \dots & \dots \\ -\frac{x_n x_1}{\Delta} & -\frac{x_n x_2}{\Delta} & \dots & 1 - \frac{x_n^2}{\Delta} \end{bmatrix}$$

It is not difficult to show that $|H_x - \lambda I| = 0$ has n positive eigen values, i.e., the metric h_x has signature n and hence is a positive definite.

Similar in the case $\alpha < 0$, the induced Riemannian metric will be negative definite.

To calculate the curvature we consider first the case $n = 2$. In this case the covariant and contravariant components of the form h are:

$$\begin{aligned} h_{11} &= 1 - \frac{x_1^2}{\Delta}, \quad h_{12} = h_{21} = -\frac{x_1 x_2}{\Delta}, \quad h_{22} = 1 - \frac{x_2^2}{\Delta}, \\ h^{11} &= \frac{\alpha + x_2^2}{\alpha}, \quad h^{12} = h^{21} = \frac{x_1 x_2}{\alpha}, \quad h^{22} = \frac{\alpha + x_1^2}{\alpha} \end{aligned}$$

Also the affine connections Γ_{ab}^c have the components:

$$\begin{aligned} \Gamma_{11}^1 &= -\frac{x_1(\alpha + x_2^2)}{\alpha \Delta}, \quad \Gamma_{11}^2 = -\frac{x_2(\alpha + x_2^2)}{\alpha \Delta}, \\ \Gamma_{12}^1 &= \Gamma_{21}^1 = \frac{x_2^2 x_1}{\alpha \Delta}, \quad \Gamma_{12}^2 = \Gamma_{21}^2 = \frac{x_1 x_2^2}{\alpha \Delta}, \end{aligned}$$

$$\Gamma^1_{22} = - \frac{x_1 (\alpha + x_1^2)}{\alpha \Delta} , \quad \Gamma^2_{22} = - \frac{x_2 (\alpha + x_1^2)}{\alpha \Delta} .$$

By using the formula,

$$R_{ab} = \frac{\partial \Gamma^c_{ab}}{\partial x^c} - \frac{\partial \Gamma^c_{ac}}{\partial x^b} + \Gamma^d_{ab} \Gamma^c_{dc} - \Gamma^d_{ac} \Gamma^c_{db},$$

for the Ricci tensors, it is easy to check that:

$$R_{11} = - \frac{(\alpha + x_1^2)}{\alpha \Delta} , \quad R_{12} = \frac{x_1 x_2}{\alpha \Delta} = R_{21}, \quad R_{22} = - \frac{(\alpha + x_1^2)}{\alpha \Delta} ,$$

and hence the curvature of S_+ will be

$$R = - 2 / \alpha .$$

Now, for $n = 3$, the Ricci tensors are given by:

$$\begin{aligned} R_{11} &= - \frac{2}{\alpha \Delta} (\alpha + x_2^2 + x_3^2) , & R_{12} &= R_{21} = \frac{2x_1x_2}{\alpha \Delta} \\ R_{22} &= - \frac{2}{\alpha \Delta} (\alpha + x_1^2 + x_3^2) , & R_{23} &= R_{32} = \frac{2x_2x_3}{\alpha \Delta} , \\ R_{33} &= - \frac{2}{\alpha \Delta} (\alpha + x_1^2 + x_2^2) , & R_{13} &= R_{31} = \frac{2x_1x_3}{\alpha \Delta} , \end{aligned}$$

and hence $R = - \frac{(3)(2)}{\alpha}$. This can be generalized for the n case

for the hypersurfaces $S(\alpha)$ where $\alpha > 0$ and once again for $\alpha < 0$ to have

$$R = \begin{cases} n(n-1) / \alpha , & n \geq 2 \\ 1 / \alpha , & n = 1 , \end{cases}$$

where the curvature R is negative for $\alpha > 0$ and positive for $\alpha < 0$.

REFERENCES

HAWKINS, S.W. (1973). *The Large Scale Structure of Space - Time*, Cambridge University Press.

WOLF, J.A. (1967). *Spaces of Constant Curvature*. McGraw - Hill, New York,