

# COMMUNICATIONS

DE LA FACULTÉ DES SCIENCES  
DE L'UNIVERSITÉ D'ANKARA

Série A: Mathématiques, Physique et Astronomie

---

TOME 22 A

ANNÉE 1973

---

## **The Mössbauer Effect**

by

**Mehmet AYDIN**

**6**

Faculté des Sciences de l'Université d'Ankara  
Ankara, Turquie

# Communications de la Faculté des Sciences de l'Université d'Ankara

Comité de Rédaction de la Série A

C. Uluçay, E. Erdik, N. Doğan

Secrétaire de publication

N. Gündüz

---

La Revue "Communications de la Faculté des Sciences de l'Université d'Ankara" est un organe de publication englobant toutes les disciplines scientifiques représentées à la Faculté: Mathématiques pures et appliquées, Astronomie, Physique et Chimie théorique, expérimentale et technique, Géologie, Botanique et Zoologie.

La Revue, à l'exception des tomes I, II, III, comprend trois séries

Série A: Mathématiques, Physique et Astronomie.

Série B: Chimie.

Série C: Sciences naturelles.

En principe, la Revue est réservée aux mémoires originaux des membres de la Faculté. Elle accepte cependant, dans la mesure de la place disponible, les communications des auteurs étrangers. Les langues allemande, anglaise et française sont admises indifféremment. Les articles devront être accompagnés d'un bref sommaire en langue turque.

Adres: Fen Fakültesi Tebliğler Dergisi Fen Fakültesi, Ankara, Turquie.

# The Mössbauer Effect

Mehmet AYDIN

*Department of Metallurgical Engineering  
Faculty of Engineering  
Middle East Technical University*

## ABSTRACT

The essence of the Mössbauer effect is not  $\gamma$ -ray emission without recoil, but it is  $\gamma$ -ray emission without transferring any nuclear transition energy to the internal energy of the crystalline solid. The problem has been first treated classically by starting with the kinematics of  $\gamma$ -ray emission for an idealized nucleus-plus-lattice system. Next, using the basic ideas of quantum mechanics, it has shown that the effect is a quantum mechanical phenomenon and can not be explained classically.

## 1. INTRODUCTION

Mössbauer effect is the phenomenon of recoil-free emission and resonant absorption of nuclear gamma rays emitted from atomic nuclei chemically bound in a crystalline solid. It allows precise determination of extremely small energy shifts as a result of the Doppler effect modulation of gamma-ray energy. The effect was discovered by Rudolph L. Mössbauer in 1957. The Physics of the Mössbauer effect has already been discussed by many authors [1-11]. The purpose of this paper is to discuss briefly the basic features of this phenomenon as clearly as possible and present it in an easy way to understand.

When a free nucleus in an excited state decays to its ground state with the emission of a  $\gamma$ -ray, then the atom normally recoils and the energy of the emitting gamma ray will depend on the kinetic energy of the recoiled atom. The laws of conservation of energy and momentum requires that the recoil energy of the nucleus of mass  $m$  upon the emission of a gamma ray of frequency  $\nu$  is

$$(1) \quad E_r = \frac{(h\nu)^2}{2 mc^2}$$

and the energy of the  $\gamma$ -ray would be  $E_0 - E_r$  where  $E_0$  is the nuclear transition energy.

Because of its considerably high energy, generally, a  $\gamma$ -photon imparts a measurable recoil to the free nucleus during emission, so that part of the excitation energy of the nucleus is lost as recoil energy. Hence any  $\gamma$ -photon emitted has insufficient energy to excite a nucleus of an absorber.

If the atom is tightly bound to the lattice, the entire solid rather than the individual atom recoils and the  $\gamma$ -ray might have the excitation energy  $E_0$  of the nucleus. Thus, we must emphasize here that the Mössbauer effect is not  $\gamma$ -ray emission without recoil, but rather is  $\gamma$ -ray emission without transfer of energy to internal degrees of freedom of the lattice and the subsequent inverse process which is called recoil-free resonant absorption and the reemission of gamma ray. The following is the classical and quantum mechanical discussions of  $\gamma$ -ray emission under certain condition giving Mössbauer effect.

## 2. CLASSICAL DISCUSSION OF MÖSSBAUER EFFECT

Consider the kinematics of  $\gamma$ -ray emission from atomic nuclei bound in a crystalline solid. Let the mass of the emitting nucleus be  $m_1$  and that of the remaining part of the solid be  $m_2$  ( $m_2 \gg m_1$ ). We assume that  $m_1$  and  $m_2$  are bound to each other by a massless spring of force constant  $k$  and unstretched length  $a$ . It makes an idealized nucleus-plus-lattice system which can be schematized as shown in Fig. (1). This is just a simplified




Figure 1. The Einstein Model of a Lattice

Einstein model of a lattice and it is an oscillator of single vibrational mode of frequency  $\omega = (k/\mu)^{1/2}$  where  $\mu$  is the reduced mass of the system.

Imagine that the nucleus of mass  $m_1$  emits a gamma ray in the transition between the excited state of energy  $E_0$  to the ground state in the direction of x-axis. Before emission the system consists of the two masses  $m_1$  and  $m_2$  at rest with the nucleus in energy state  $E_0$ . After emission it consists of two masses  $m_1$  and  $m_2$  with velocities  $\dot{x}_1$  and  $\dot{x}_2$ , respectively, with the nucleus in ground state and an emitted gamma photon of frequency  $\nu$ . The equations of momentum and energy balance are then,

$$\text{Momentum} : \frac{h\nu}{c} = m_1\dot{x}_1 + m_2\dot{x}_2$$

$$\text{Energy} : E_0 = h\nu + \frac{1}{2} m_1 \dot{x}_1^2 + \frac{1}{2} m_2 \dot{x}_2^2 + \frac{1}{2} k (x_2 - x_1 - a)^2$$

These equations become more useful when they are expressed in terms of relative and center of mass coordinates  $x$  and  $u$ , respectively, defined as

$$x = x_2 - x_1$$

$$u = \frac{m_1 x_1 + m_2 x_2}{m_1 + m_2}$$

Here a little algebra shows that the above equations become

$$(2) \quad M\dot{u} - \frac{h\nu}{c} = 0$$

$$(3) \quad E_0 = h\nu + \frac{1}{2} M \dot{u}^2 + \frac{1}{2} \mu \dot{x}^2 + \frac{1}{2} k (x - a)^2$$

where  $M$  is the total mass ( $M = m_1 + m_2$ ). From Eqs. (2) and (3) we get the following equation

$$(4) \quad E_0 = h\nu + \frac{1}{2} \frac{(h\nu)^2}{Mc^2} + E_{int}$$

where  $E_{int} = \frac{1}{2} \mu \dot{x}^2 + \frac{1}{2} k (x-a)^2$

As we see,  $E_{int}$  depends only on the relative coordinate  $x$  and called the "internal energy" of the system. A comparison of the

Eqs. (1) and (4) shows that the second term on the right hand side of the Eq. (4) is what we previously called the "recoil energy". The only difference is that the total mass  $M$  and not  $m_1$  appears in this term. Since  $M$  is enormously large, the recoil energy is negligible and to a good approximation,

$$(5) \quad h\nu \cong E_0 - E_{int}$$

The Mössbauer effect begins to emerge in this equation. Here, if it were possible for  $E_{int}$  to be zero, we would have

$$(6) \quad h\nu \cong E_0$$

which is just the mathematical statement of the Mössbauer effect. It describes a transition between an excited state of energy  $E_0$  to the ground state, in which the total energy released in the transition is all given to the gamma photon. In other words, if  $E_{int}$  can be equal to zero under certain condition, or in other language, if no internal lattice vibrations (phonons) are excited, we have the possibility of the Mössbauer effect.

The question is now under which circumstances  $E_{int}$  can be equal to zero. In classical mechanics it is quite clear that it can not be so. Upon the emission of gamma ray, the individual atom suffers a sudden momentum impulse and therefore it transfers a finite momentum to the system which make the internal energy of the lattice to be changed. But, from quantum mechanical point of view the story is different and one can transfer a momentum to a nucleus without changing the internal energy of the system [9].

### 3. QUANTUM MECHANICAL DISCUSSION OF MÖSSBAUER EFFECT

For simplicity, we consider again the Einstein lattice of Fig. (1). Quantum mechanically, the internal energy of this system is quantized and the allowed values are just the oscillator eigenvalues  $E_{int} = (n + 1/2)\hbar w$ . For the system, say initially in the ground state ( $n = 0$ ), if there is a finite probability to be still in the ground state after gamma ray emission, the Mössbauer effect will be possible. The purpose now is to calculate this probability called "Debye-Waller factor" and show that it is different from zero.

Let  $P_{oo}$  be the probability amplitude for the system to remain in this original oscillator ground state after emitting a gamma photon of momentum  $p_o = \frac{h\nu}{c}$ . The probability amplitude for the nucleus to be in the oscillator ground state, when it has momentum  $p$ , is just the ground state wave function  $\Phi_o(p)$ . Since the nucleus was in the ground state before recoil and it gets an additional momentum  $p_o$  after the emission of gamma photon, the probability amplitude for the nucleus to have momentum  $p$  after recoil is the same as the probability amplitude for the nucleus to have had momentum  $p-p_o$  before recoil and it is just  $\Phi_o(p-p_o)$ . Therefore, the probability amplitude  $P_{oo}$  can now be written as [12]

$$(7) \quad P_{oo} = \int \Phi_o^*(p) \cdot \Phi_o(p-p_o) dp$$

The asterisk\* on the right hand side stands for complex conjugate.

In order to evaluate the above integral, we introduce the ground state space function  $\Psi_o(x)$ , of which  $\Phi_o(p)$  is the Fourier transform

$$(8) \quad \Phi_o(p) = \frac{1}{\sqrt{2\pi\hbar}} \int \Psi_o(x) e^{-ipx/\hbar} dx$$

If we put this into Eq. (7), we obtain

$$(9) \quad P_{oo} = \frac{1}{2\pi\hbar} \iiint \Psi_o^*(x') e^{ipx'/\hbar} \Psi_o(x) e^{-i(p-p_o)x/\hbar} dp dx dx'$$

Substituting the delta function into Eq. (9), we obt in

$$(10) \quad P_{oo} = \iint \Psi_o^*(x') \delta(x-x') \cdot \Psi_o(x) e^{ipox/\hbar} dx dx'$$

Remembering that  $\Psi_o(x) = \int \Psi_o(x) \delta(x-x') dx'$ , then Eq. (10) becomes

$$(11) \quad P_{oo} = \int \Psi_o^*(x) \Psi_o(x) e^{ipox/\hbar} dx$$

We can now find the probability  $f$  for a transition without energy loss which is

---


$$\hbar = \frac{h}{2\pi}$$

$$(12) \quad |P_{00}|^2 = f = \left| \int \Psi_0^*(x) \Psi_0(x) e^{ip_0x/\hbar} dx \right|^2$$

The ground state space function for our system which is a harmonic oscillator is given as

$$\Psi_0(x) = \sqrt{\frac{mw}{\pi \hbar}} e^{-\frac{mw}{2\hbar} x^2}$$

Where  $m$  is the mass of the nucleus. Substituting this into Eq. (12), we finally obtain

$$(13) \quad f = e^{-\frac{E_r}{\hbar w}}$$

where  $E_r$  is the recoil energy of the free nucleus defined by Eq. (1).

As we see, for the Einstein solid the problem is quite simple. If the energy  $E_r$  is of the same order as  $\hbar w$ , the quantized level spacing of the oscillator, the probability for the Mössbauer effect will be substantial. When the recoil energy is small compared to the smallest amount of energy ( $E_e = \hbar w_e$ ) that can be given to the solid the lattice will not be excited and the  $\gamma$ -ray will have the total transition energy.

Eq. (13) shows that, for a given nucleus of mass  $m$  and nuclear transition energy  $E_0$ , if  $E_r$  is large compared to  $\hbar w$  then upon the emission of each  $\gamma$ -photon the lattice will be excited and the probability of obtaining the Mössbauer effect will be small.

We must remember that here we worked out a rather idealized problem in which all possible degrees of freedom and all frequencies of the original problem were replaced by a single degree of freedom and a single frequency. In general, one can use the more accurate model of a lattice, such as the Debye model, and obtain a similar result [6]

$$f = e^{-\frac{3}{2} \frac{E_r}{\hbar w_d}}$$

where  $w_d$  is the Debye frequency, and  $E_d = \hbar w_d$  is the maximum lattice vibration energy.



## REFERENCES

- [ 1 ] R. L. Mössbauer, Z. Physik, 151, 124 (1958).
- [ 2 ] R. L. Mössbauer and D. H. Sharp, Rev. Mod. Phys. 36, 410 (1964).
- [ 3 ] H. Lustig, Am. J. Phys. 29, 1 (1961).
- [ 4 ] W. E. Burcham, F. R. S., Science Progress, 630 (1960)
- [ 5 ] G. K. Wertheim, phys. Rev. Letters, 4, 403 (1960).
- [ 6 ] H. Fraunfelder, The Mössbauer Effect, W. A. Benjamin, N. Y. 1962, PP. 23
- [ 7 ] L. J. Eyges, AFCRL - 64 - 840.
- [ 8 ] E. Fluck, W. Kerler, W. Neuwirth, Angew. Chem. 2, 277 (1963).
- [ 9 ] H. Lipkin, Ann. phys. (n. Y.), 18, 182 (1962).
- [10] D. A. O 'Connor, Contemp, phys. 9, 521 (1968).
- [11] W. M. Visscher, Ann. phys. (N. Y.), 9, 194 (1960).
- [12] R. P. Feynmann, "Lecture on physics", Vol. III, Addison-wesley, New York, 1965, PP. 5-12.

## ÖZET

Mössbauer olayı ahlnda  $\gamma$ -ışınlarının geri tepmesiz olarak neşredilmesi olayı değildir; fakat, çekirdek dönüşüm enerjisinden kristalin iç enerjisine hiç enerji transferi olmadan  $\gamma$ -ışımın neşri olayıdır. Einstein kistal modeli esas alınarak, uyarılmış çekirdekten  $\gamma$ -ışımın neşri olayı önce klasik olarak incelenmiş ve kristalin iç enerjisinde bir değişme yapmadan bir  $\gamma$ -photonunun neşrolunamayacağı görülmüştür. Kuantum mekaniği açısından durum farklıdır ve bazı şartlarda bir  $\gamma$ -ışımın, çekirdek enerjisine eşit bir enerji ile neşrolunması mümkündür.

**Prix de l'abonnement annuel**

Turquie : 15 TL; Étranger: 30 TL.

Prix de ce numéro : 5 TL (pour la vente en Turquie).

Prière de s'adresser pour l'abonnement à : Fen Fakültesi

Dekanlığı Ankara, Turquie.